





## 1-(Arenesulfonyloxy) benziiodoxolones



Treatment of aromatic compounds with 1-(arenesulfonyloxy) benziiodoxolones and iodine gave the corresponding iodinated compounds in good yield.

Using a halide salt such as lithium bromide or lithium chloride instead of iodine, the corresponding aryl bromide and chloride were also obtained in good yields <sup>1,2</sup>.



- 1) Muraki, T.; Togo, H.; Yokoyama, M. *Synlett* 1998(3), 286-288
- 2) Muraki, T.; Togo, H.; Yokoyama, M. *J. Org. Chem.* 1999, 64, 2883-2889
- 3) Alkynyliodonium Salts in Organic Synthesis. Zhdankin, V.V.; Stang, P.J. *Tetrahedron* 1998, 54, 10927-10966.
- 4) IBX Amides: New Family of Hypervalent Iodine Reagents. Zhdankin, V.V.; Kuposov, A.Y.; Netzel, B.C.; Yashin, N.V.; Rempel, B.P.; Ferguson, M.J.; Tykwinski, R.R. *Angew. Chem. Int. Ed.* 2003, 42, 2194-2196.
- 5) Synthesis and Structure of Amino Acid-Derived Benziiodazoles: New Hypervalent Iodine Heterocycles. Zhdankin, V.V.; Kuposov, A.Y.; Su, L.; Boyarskikh, V.V.; Netzel, B.C.; Young, V.G. *Org. Lett.* 2003, 5, 1583-1586
- 6) Zhdankin, V.V.; Kuehl K.J.; Krasusky A.P.; Boltz J.T.; Simonsen A.J. *J. Org. Chem.* 1996 (61) 6547-6551